arXiv:1410.5638v1 [math.DG] 21 Oct 2014

A GENERALIZED PALAIS-SMALE CONDITION IN THE FRÉCHET SPACE SETTING

KAVEH EFTEKHARINASAB

In memory of Vladimir Sharko

Abstract. We extend the Palais-Smale condition to the cases of C^1-functionals on Fréchet spaces and Fréchet-Finsler manifolds of class C^1. In these contexts we use the introduced conditions along with Ekeland's variational principle to obtain some customary results concerning the existence of minima.

1. Introduction

The Palais-Smale condition is an essential tool for critical point theory in infinite dimensional spaces. Since it was established by Palais and Smale [8], a number of its generalizations for various categories of maps have been introduced (cf. [6] for a survey). The Palais-Smale type conditions have been applied to numerous problems in critical points theory and its applications to various problems in differential equations, geometry and physics (see [6]).

In this paper we generalize the Palais-Smale condition to the cases of C^1-functionals on Fréchet spaces and Fréchet-Finsler manifolds of class C^1. It is instantly noticed that due to the weak structural constraints of Fréchet spaces, applications of the conditions are too restrictive. It is well known that some of the basic tools for locating critical points of C^1-functionals are the deformation lemmas. The deformations that appear in the deformation lemmas are obtained as solutions of Cauchy problems. But on Fréchet spaces, an ordinary differential equation may admit no, one or multiple solutions for the same initial condition. Thus, we can not get deformation results for Fréchet spaces in general. However, we can still use the introduced conditions along with Ekeland's variational principle to locate critical points. We then apply them and the works of Ekeland [2] and Qui [9] to state some standard results dealing with the existence of minima.

2010 *Mathematics Subject Classification.* 58E05, 58E30, 58B20.

Key words and phrases. The Palais-Smale condition, Fréchet-Finsler manifold, Ekelend's variational principle.

Preprint submitted to Fundamenta Mathematicae

2. Prerequisites

We use the following notion of C^k-maps in Michal-Bastiani sense. We shall work in the category of manifolds of class at least C^1.

Definition 2.1. *Let E and F be Fréchet spaces, $U \subseteq E$ open and $f : U \to F$ a continuous map. The derivative of f at $x \in U$ in the direction of $h \in E$ is defined as*

$$\mathrm{d}\, f(x)(h) \coloneqq \lim_{t \to 0} \frac{1}{t}(f(x + ht) - f(x))$$

whenever the limit exits. The map f is called differentiable at x if $\mathrm{d}\, f(x)(h)$ exists for all $h \in E$. It is called a C^1-map if it is differentiable at all points of U and

$$\mathrm{d}\, f : U \times E \to F, \quad (x, h) \mapsto \mathrm{d}\, f(x)(h)$$

is a continuous map. The map f is called a C^k-map, $k \geqslant 2$, if it is continuous, the higher directional derivatives

$$\mathrm{d}^k f(x)(h_1, \cdots, h_k) \coloneqq \lim_{t \to 0} \frac{1}{t}(\mathrm{d}^{k-1} f(x + h_k t)(h_1, \cdots, h_{k-1}) - \mathrm{d}^{k-1} f(x)(h_1, \cdots, h_{k-1}))$$

exist for all integers $j \leqq k$, $x \in U$ and $h_1, \cdots, h_j \in E$, and all maps

$$\mathrm{d}^j f : U \times E^j \to F, \quad (x, h_1, \cdots, h_j) \mapsto \mathrm{d}^j f(x)(h_1, \cdots, h_j)$$

are continuous. We say that f is smooth or C^∞ if it is C^k for all $k \in \mathbb{N}$.

Within this framework most of the notions of the theory of infinite dimensional manifolds are sufficiently well behaved for our purposes. In particular, Fréchet manifolds, Fréchet tangent bundles and C^k-maps between Fréchet manifolds are defined in obvious way (cf. [3]). However, for a Fréchet manifold M modelled on a Fréchet space F we can define the set-theoretic cotangent bundle $T'M \coloneqq \bigcup_{m \in M} (T_m M)'$, where $(T_m M)'$ is the dual of a tangent space $T_m M$. As a set, it carries a natural structure of a vector bundle over M, but in general there is no vector topology on F', the dual of F, that can lead to the identification $T'M \cong F \times F'$, see [7, Remark I.3.9]. Thus instead, we use the notion of a $\mathcal{B}$-cotangent bundle [10]. In this definition to put a manifold structure on $T'M$, the dual of F is equipped by a $\mathcal{B}$-topology, where $\mathcal{B}$ is a bornology on F. To be precise, we recall that a family $\mathcal{B}$ of bounded subsets of F that covers F is called a bornology on F if it is directed upwards by inclusion and if for every $B \in \mathcal{B}$ and $r \in \mathbb{R}$ there is a $C \in \mathcal{B}$ such that $r \cdot B \subset C$.

Let E be a Fréchet space, $\mathcal{B}$ a bornology on E and $L_{\mathcal{B}}(E, F)$ the space of all linear continuous maps from E to F. The $\mathcal{B}$-topology on $L_{\mathcal{B}}(E, F)$ is a Hausdorff locally convex topology defined by all seminorms obtained as follows:

$$\mathrm{P}^n_B(L) \coloneqq \sup\{\mathrm{p}_n(L(e)) \mid e \in B\}, \tag{2.1}$$

where $B \in \mathcal{B}$ and $\{\mathrm{p}_n\}_{n\in\mathbb{N}}$ is a family of seminorms defining the topology of F. One similarly may define $L^k_{\mathcal{B}}(E,F)$ and $\bigwedge^k L_{\mathcal{B}}(E,\mathbb{R})$, the space of k-linear jointly continuous maps of E^k to F and the space of anti-symmetric k-linear jointly continuous maps of E^k to $\mathbb{R}$, respectively. If $\mathcal{B}$ contains all compact sets, then the $\mathcal{B}$-topology on the space $L_{\mathcal{B}}(E,\mathbb{R}) = E'_{\mathcal{B}}$ of all continuous linear functional on E, the dual of E, is the topology of compact convergence. For this and further details on bornologies we refer to [4].

If $\mathcal{B}$ contains all compact sets of E, then $L^k_{\mathcal{B}}(E, L^l_{\mathcal{B}}(E,F))$ is canonically isomorphic to $L^{l+k}_{\mathcal{B}}(E,F)$ as a topological vector space, see [5, Theorem 0.1.3]. In particular, $L^2_{\mathcal{B}}(E,R) = L^2_{\mathcal{B}}(E) \cong L_{\mathcal{B}}(E, E'_{\mathcal{B}})$. Under the above condition on $\mathcal{B}$, we define the differentiability of class $C^k_{\mathcal{B}}$: Let $U \subset E$ be open, a map $f : U \to F$ is called $C^1_{\mathcal{B}}$ if its partial derivatives exist and the induced map $\mathrm{d}\, f : U \to L_{\mathcal{B}}(E,F)$ is continuous. Similarly we can define maps of class $C^k_{\mathcal{B}}$, $k \in \mathbb{N} \cup \{\infty\}$, see [5, Definition 2.5.0]. A map $f : U \to F$ is $C^k_{\mathcal{B}}, k \geqq 1$, if and only if f is C^k in the sense of Definition (2.1), see [5, Theorem 2.7.0 and Corollary 1.0.4 (2)]. In particular, f is $C^\infty_{\mathcal{B}}$ if and only in f is C^∞. Thus, if f at $x \in F$ is C^1 and hence $C^1_{\mathcal{B}}$, the derivative f at x, $\mathrm{d}\, f(x)$, is an element of $E'_{\mathcal{B}}$, and we shall denote $\mathrm{d}\, f(x)$ evaluated at $h \in F$ by $\mathrm{d}\, f(x)(h)$.

Assume that $\mathcal{B}$ is a bornology on F containing all compact sets and M is a Fréchet manifold modelled on F. Let f be a functional defined over M. The derivative of f at $x \in M$ can be written in terms of the iterated tangent bundles of M and we can consider $\mathrm{d}\, f : TM \to F$ given by $\mathrm{d}\, f(x,h) = \mathrm{d}\, f(x)(h)$ upon locally identifying TM with $U \times F$, where U is an open set in F. Therefore, if at $x \in M$ a map $f : M \to \mathbb{R}$ is C^1 and hence $C^1_{\mathcal{B}}$, then $\mathrm{d}\, f(x)$ belongs to $L_{\mathcal{B}}(T_xM, \mathbb{R}) = (T_xM)'_{\mathcal{B}}$.

Definition 2.2. *Let M be a Fréchet manifold modelled on a Fréchet space F and $\mathcal{B}$ a bornology on F. The $\mathcal{B}$-cotangent bundle of M is defined as $TM'_{\mathcal{B}} \coloneqq \bigcup_{x\in M}(T_xM)'_{\mathcal{B}}$ and the k-exterior product of the $\mathcal{B}$-cotangent bundle as $\bigwedge^k TM'_{\mathcal{B}} \coloneqq \bigcup_{x\in M} \bigwedge^k (T_xM)'_{\mathcal{B}}$.*

If $\mathcal{B}$ is chosen such that $T(\mathcal{B}) \subset \mathcal{B}$ for all continuous linear endomorphisms T of F, then $\bigwedge^k TM'_{\mathcal{B}}$ is a vector bundle in the category of locally convex spaces with the local model $F \times \bigwedge^k F'_{\mathcal{B}}$. In particular, $TM'_{\mathcal{B}}$ is a vector bundle in the category of locally convex spaces with the local model $F \times F'_{\mathcal{B}}$, see [10, Remark (1), p. 339].

3. The Palais-Smale condition

A Finsler structure for a Fréchet manifold M in the sense of Palais was defined in [1]. We now define a Finsler structure on its $\mathcal{B}$-cotangent bundle.

Henceforth, we always assume that a bornology $\mathcal{B}$ on a Fréchet space F contains all compact sets and we always chose $\mathcal{B}$ such that $T(\mathcal{B}) \subset \mathcal{B}$ for all continuous linear endomorphisms T of F whenever we define the $\mathcal{B}$-cotangent bundle of a manifold M modelled on F.

Definition 3.1. *Let M be a C^k Fréchet manifold modelled on a Fréchet space F, where $k \in \mathbb{N} \cup \{\infty\}$. A Finsler structure for M is a collection of functions $\| \cdot \|^n_M \colon TM \to \mathbb{R}^+$, $n \in \mathbb{N}$, on its tangent bundle TM such that*

(1) *For every $x \in M$, $\| \cdot \|^n_x := \| \cdot \|^n_{M|_{T_xM}} \colon T_xM \to \mathbb{R}^+$ is a collection of seminorms on the tangent space T_xM such that for every chart $\varphi : U \to F$ with $x \in U$, the collection of seminorms $v \in F \mapsto \| \, \mathrm{d}\, \varphi^{-1}(\varphi(x))(v) \, \|_x$ generates the topology of F.*

(2) *Given $K > 1$, $x_0 \in M$ and a chart $\varphi : U \to F$ with $x_0 \in U$, there exists an open neighborhood $\mathcal{U}$ of x_0 in U such that*

$$\frac{1}{K} \| \, \mathrm{d}\, \varphi^{-1}(\varphi(x_0))(v) \, \|^n_{x_0} \leqq \| \, \mathrm{d}\, \varphi^{-1}(\varphi(x))(v) \, \|^n_x \leqq \ K \| \, \mathrm{d}\, \varphi^{-1}(\varphi(x_0))(v) \, \|^n_{x_0}$$

for all $x \in \mathcal{U}$, $n \in \mathbb{N}$, $v \in F$.

If $\{\| \cdot \|^n\}_{n\in\mathbb{N}}$ is a Finsler structure on TM then eventually we can obtain a graded Finsler structure, denoted by $(\| \cdot \|^n)_{n\in\mathbb{N}}$, on TM. A Fréchet manifold M is called Fréchet-Finsler if its tangent bundle TM admits a Finsler structure. If a Fréchet manifold M is paracompact and its model space F is such that all seminorms defining the topology of F are smooth maps on $F\backslash\{0\}$, then TM admits a Finsler structure ([1, Proposition 4]).

Let M be a connected Fréchet-Finsler manifold endowed with a graded Finsler structure $(\| \cdot \|^n)_{n\in\mathbb{N}}$. For all $n \in \mathbb{N}$, define $d^n(x, y) = \inf_\gamma L^n(\gamma)$, where infimum is taken over all C^1-curve $\gamma : [a, b] \to M$ such that $\gamma(a) = x$, $\gamma(b) = y$ and

$$L^n(\gamma) = \int_a^b \| \, (\gamma'(t)) \, \|^n_{\gamma(t)} \, \mathrm{d}t.$$

Therefore, we obtain an increasing sequence of pseudometric $(d^n(x, y))_{n\in\mathbb{N}}$. Define

$$\rho(x, y) = \sum_{n=1}^{n=\infty} \frac{1}{2^n} \frac{d^n(x, y)}{1 + d^n(x, y)}. \tag{3.1}$$

This is a metric, called the Finsler metric, consistent with the original topology of M ([1]). In the sequel, we say that M is complete if M is complete with respect to this metric. We say that ρ is locally compatible if for each $p \in M$ there exist a chart $(p \in U, \varphi)$ and constants α, β such that for all $n \in \mathbb{N}$ and $x, y \in U$

$$\alpha\rho(x, y) \leqq \mathrm{p}_n(\varphi(x) - \varphi(y)) \leqq \beta\rho(x, y),$$

where $\{\mathrm{p}_n\}_{n\in\mathbb{N}}$ is a family of seminorms defining the topology of F.

The $\mathcal{B}$-cotangent bundle $TM'_{\mathcal{B}}$ of M carries a natural Finsler structure, denoted again by $\{\|\cdot\|^n\}_{n\in\mathbb{N}}$, characterized by letting

$$\| w \|^n_x = \sup\{< w, v > | \; v \in T_xM, \| v \|^n_x = 1\},$$

where $x \in M$, $w \in (T_xM)'_{\mathcal{B}}$ and $< \cdot, \cdot >$ is the duality pairing between T_xM and $(T_xM)'_{\mathcal{B}}$ which is separately continuous. Consequently, for a C^1-functional $f : M \to \mathbb{R}$ the mappings $\| \mathrm{d}\, f \|^n_x \colon M \to \mathbb{R}$ given by $x \mapsto \| \mathrm{d}\, f(x) \|^n_x$ are well defined and continuous for all $n \in \mathbb{N}$. We shall omit the index x in $\| \mathrm{d}\, f(x) \|^n_x$ and denote it just by $\| \mathrm{d}\, f(x) \|^n$.

Definition 3.2. *Let F be a Fréchet space, $\mathcal{B}$ a bornology on F and $F'_{\mathcal{B}}$ the dual of F equipped with the $\mathcal{B}$-topology. Let $f : F \to \mathbb{R}$ be a C^1-functional.*

(i) *We say that f satisfies the Palais-Smale condition,* (PS) *in short, if each sequence $(x_i) \subset F$ such that $f(x_i)$ is bounded and*

$$\mathrm{d}\, f(x_i) \to 0 \quad \text{in} \quad F'_{\mathcal{B}},$$

has a convergent subsequence.

(ii) *We say that f satisfies the Palais-Smale condition at level $c \in \mathbb{R}$,* $(\mathrm{PS})_c$ *in short, if each sequence $(x_i) \subset F$ such that*

$$f(x_i) \to c \quad \text{and} \quad \mathrm{d}\, f(x_i) \to 0 \quad \text{in} \quad F'_{\mathcal{B}},$$

has a convergent subsequence.

Definition 3.3. *Let M be a connected Fréchet-Finsler manifold endowed with a Finsler structure $\{\|\cdot\|^n\}_{n\in\mathbb{N}}$ and the corresponding Finsler metric ρ. Let $f : M \to \mathbb{R}$ be a C^1-functional.*

(i) *We say that f satisfies the Palais-Smale condition,* (PS) *in short, if for all $n \in \mathbb{N}$ each sequence $(x_i) \subset M$ such that $f(x_i)$ is bounded and $\| \mathrm{d}\, f(x_i) \|^n \to 0$, has a convergent subsequence.*

(ii) *We say that f satisfies the Palais-Smale condition at level $c \in \mathbb{R}$,* $(\mathrm{PS})_{\mathrm{c}}$ *in short, if for all $n \in \mathbb{N}$ each sequence $(x_i) \subset M$ such that $f(x_i) \to c$ and $\| \mathrm{d}\, f(x_i) \|^n \to 0$, has a convergent subsequence.*

Let f be a C^1-functional on a Fréchet space (or a Fréchet-Finsler manifold). As usual, a point p in the domain of f is said to be a critical point of f if $\mathrm{d}\, f(p) = 0$, the corresponding value $c = f(p)$ will be called a critical value. We use the following standard notation: $K_c \coloneqq \{p \in M \mid f(p) = c, \mathrm{d}\, f(p) = 0\}$.

4. Existence of minima

Consider the following particular version of Ekeland's variational principle (cf. [2]).

Theorem 4.1. *Let (X, σ) be a complete metric space. Let a functional $f : X \to (-\infty, \infty]$ be semi-continuous, bounded from below and not identical to $+\infty$. Let $a > 1$, $b > 0$ and $x \in X$ be given such that $f(x) \leqq \inf_X f + a$. Then there exists $x_a \in M$ such that*

(1) $f(x_a) \leqq f(x)$,
(2) $\sigma(x_a, x) \leqq \dfrac{1}{b}$,
(3) $f(x_a) < f(x) + ab\sigma(x, x_a), \quad \forall x \neq x_a \in M.$

When X is a complete Fréchet-Finsler manifold and f is of class C^1, we have the following:

Theorem 4.2. *Let M be a connected complete Fréchet-Finsler manifold of class C^1 endowed with a Finsler structure $\{\| \cdot \|^n\}_{n\in\mathbb{N}}$ and the corresponding Finsler metric ρ which is locally compatible. Let a C^1-functional $f : M \to \mathbb{R}$ be bounded from below. Then, for each $\theta > 1$ and each $m \in M$ such that $f(m) \leqq \inf_M f + \theta^2$, there exist $m_\theta \in M$ and a constant ε_θ such that*

(1) $f(m_\theta) \leqq f(m)$,
(2) $\rho(m_\theta, m) \leqq \theta$,
(3) $\| \mathrm{d} f(m_\theta) \|^n \leqq \varepsilon_\theta, \quad \forall n \in \mathbb{N}.$

Proof. Given $\theta > 1$, apply Theorem 4.1 with $b = \dfrac{1}{\theta}$ to find $m_\theta \in M$ such that (1) and (2) hold and

$$f(x) > f(m_\theta) - \theta\rho(m_\theta, x), \quad \forall x \neq m_\theta \in M. \tag{4.1}$$

Since ρ is locally compatible there exist a chart $\varphi : U \to F$ with $m_\theta \in U$ (F is the local model of M) and constants α, β such that for all $n \in \mathbb{N}$ and $x, y \in U$

$$\alpha\rho(x, y) \leqq \mathrm{p}_n(\varphi(x) - \varphi(y)) \leqq \beta\rho(x, y), \tag{4.2}$$

where $\{\mathrm{p}_n\}_{n\in\mathbb{N}}$ is a family of seminorms defining the topology of F. Let $h \in \varphi(U)$ and $t > 0$ be such that $\varphi^{-1}(\varphi(m_a) + th) \in U$. Therefore, by (4.1)

$$f(\varphi^{-1}(\varphi(m_\theta))) - f(\varphi^{-1}(\varphi(m_\theta) + th)) < \theta\rho(m_\theta, \varphi^{-1}(\varphi(m_\theta) + th)).$$

Hence, for all $n \in \mathbb{N}$

$$\begin{aligned} f(\varphi^{-1}(\varphi(m_\theta))) - f(\varphi^{-1}(\varphi(m_\theta) + th)) &< \frac{\theta}{\alpha}\mathrm{p}_n(\varphi(m_\theta) - (\varphi(m_\theta) + th)) = \frac{\theta}{\alpha}\mathrm{p}_n(h)t \\ &< \frac{\theta}{\alpha}\beta\rho(m_\theta, \varphi^{-1}(\varphi(m_\theta) + th))t < \frac{\theta^2}{\alpha}\beta t \end{aligned} \tag{4.3}$$

Define a C^1-curve $\gamma : [0, t_0] \to M$ by $\gamma(t) = \varphi^{-1}(\varphi(m_\theta) + th)$ such that $\gamma([0, t_0]) \subset U$ and let $\varepsilon_\theta = \dfrac{\theta^2}{\alpha}\beta$. By dividing both side in (4.3) by t and letting $t \to 0$, we obtain

$$\mathrm{d}\, f(m_\theta)(h) = \frac{\mathrm{d}}{\mathrm{d}t}(f \circ \gamma) \mid_{t=0} \geqslant -\varepsilon_\theta.$$

Changing h with $-h$ gives $\mathrm{d}\, f(m_\theta)(h) \leqq \varepsilon_\theta$, thereby for all $n \in \mathbb{N}$

$$\| \mathrm{d}\, f(m_\theta) \|^n \leqq \sup_{\mathrm{p}_i(h)=1, \forall i \in \mathbb{N}} \mathrm{d}\, f(m_\theta)(h) \leqq \varepsilon_\theta.$$

□

Corollary 4.1. *Let M and f be as in Theorem 4.2. If $(\mathrm{PS})_c$ holds with $c = \inf_M f$, then f attains its minimum at a point in K_c.*

Proof. According to Theorem 4.2, if for each $i \in \mathbb{N}$ we let $\theta^2 = \dfrac{1}{i}$, then there exists $x_i \in M$ such that

$$f(x_i) \leqq c + \frac{1}{i}, \quad \| \mathrm{d}\, f(x_i) \|^n \leqq \frac{\beta}{i\alpha}, \quad \forall n \in \mathbb{N}.$$

Therefore, $f(x_i) \to c$ and $\| \mathrm{d}\, f(x_i) \|^n \to 0$ for all $n \in \mathbb{N}$. Since f satisfies $(\mathrm{PS})_c$ it follows that there exists a convergent subsequence $x_{i_j} \to x_0$. Hence, $f(x_0) = c = \inf_M f$ and $\mathrm{d}\, f(x_0) = 0$. □

The similar results can be obtained in the case of Fréchet spaces by means of the following Ekeland's variational principle for Fréchet spaces.

Theorem 4.3 ([9], Corollary 2.1)**.** *Let F be a Fréchet space with topology generated by an increasing sequence $\mathrm{p}_1(\cdot) \leqq \mathrm{p}_2(\cdot) \leqq \cdots$ of seminorms. Let a functional $f : F \to (-\infty, \infty]$ be semi-continuous, bounded from below and not identical to $+\infty$. Let $\eta > 0$ and $x_0 \in F$ be given such that $f(x_0) \leqq \inf_F f + \eta$, and let $(\lambda_n)_{n\in\mathbb{N}}$ be a sequence of positive real numbers. Then for any $i \in \mathbb{N}$, there exists $z \in F$ such that*

(1) $\lambda_j \mathrm{p}_j(z - x_0) \leqq f(x_0) - f(z)$ *for* $j = 1, \cdots, i$;
(2) $\mathrm{p}_j(z - x_0) < \dfrac{\eta}{\lambda_j}$ *for* $j = 1, \cdots, i$;
(3) *for any $x \in F, x \neq z$, there exists $m \in \mathbb{N}$ such that*

$$\lambda_m \mathrm{p}_m(x - z) + f(x) > f(z),$$

or equivalently, $\sup_n \lambda_n \mathrm{p}_n(x - z) + f(x) > f(z)$ *for any $x \in F, x \neq z$.*

Theorem 4.4. *Let F be a Fréchet space with topology generated by an increasing sequence $\mathrm{p}_1(\cdot) \leqq \mathrm{p}_2(\cdot) \leqq \cdots$ of seminorms. Let $\mathcal{B}$ a bornology on F and $\{\mathrm{P}_B^n\}_{n\in\mathbb{N}}$ the family of seminorms (defined as in (2.1)) that generates the topology of $F'_{\mathcal{B}}$. Let a C^1-functional $f : F \to \mathbb{R}$ be bounded from below. Then, for each $\varepsilon > 0$, $i \in \mathbb{N}$, $x \in F$ such that $f(x) \leqq \inf_F f + \varepsilon$, there exists $z \in F$ such that*

(1) $\mathrm{p}_j(z-x_0) \leqq \dfrac{f(x_0)-f(z)}{\sqrt{\varepsilon}}$ *for* $j=1,\cdots,i$;
(2) $\mathrm{p}_j(z-x_0) < \sqrt{\varepsilon}$ *for* $j=1,\cdots,i$;
(3) $\mathrm{P}_B^n(\mathrm{d}\, f(z)) \leqq \sqrt{\varepsilon}, \quad \forall n \in \mathbb{N}, B \in \mathcal{B}$.

Proof. Given $\varepsilon > 0$ and $i \in \mathbb{N}$, apply Theorem 4.3 with $\lambda_j = \sqrt{\varepsilon}$ for all $j \in \mathbb{N}$ and $\eta = \varepsilon$ to find $m \in \mathbb{N}$ and $z \in F$ such that (1) and (2) hold and

$$f(x) > f(z) - \sqrt{\varepsilon}\mathrm{p}_m(x-z), \quad \forall x \neq z \in F. \tag{4.4}$$

By applying (4.4) to $x = z + th$ with $t > 0$ and $h \in F$, we obtain

$$f(z+th) - f(z) > -\sqrt{\varepsilon}\mathrm{p}_m(h)t.$$

Dividing both side by t and letting $t \to 0$ yields $\mathrm{d}\, f(z)(h) > -\sqrt{\varepsilon}\mathrm{p}_m(h)$ for all $h \in F$. Changing h with $-h$ gives $\mathrm{d}\, f(m_\theta)(h) \leqq \sqrt{\varepsilon}\mathrm{p}_m(h)$ for all $h \in F$, which means for all $n \in \mathbb{N}$ and $B \in \mathcal{B}$ we have $\mathrm{P}_B^n(\mathrm{d}\, f(z)) \leqq \sqrt{\varepsilon}$. □

Corollary 4.2. *Let F and f be as in Theorem 4.4. If $(\mathrm{PS})_c$ holds with $c = \inf_M f$, then f attains its minimum at a point in K_c.*

Proof. According to Theorem 4.4, if for each $i \in \mathbb{N}$ we let $\varepsilon = \dfrac{1}{i}$, then there exists $x_i \in F$ such that

$$f(x_i) \leqq c + \frac{1}{i}, \quad \mathrm{P}_B^n(\mathrm{d}\, f(x_i)) \leqq \frac{1}{\sqrt{i}}, \quad \forall n \in \mathbb{N}, B \in \mathbb{N}.$$

Therefore, $f(x_i) \to c$ and $\mathrm{d}\, f(x_i) \to 0$ in $F'_{\mathcal{B}}$. Since f satisfies $(\mathrm{PS})_c$ it follows that there exists a convergent subsequence $x_{i_j} \to x_0$. Hence, $f(x_0) = c = \inf_M f$ and $\mathrm{d}\, f(x_0) = 0$. □

References

[1] C. L. Bejan, Finsler structures on Fréchet bundles, Proc. 3-rd Seminar on Finsler spaces, Univ. Braşov 1984, Societatea de ştiinţe Matematice Romania, Bucharest (1984), 49-54.

[2] I. Ekeland, On the variational principle, J. Math. Anal. Appl. **47** (1974), 324-353.

[3] R. S. Hamilton, The inverse function theorem of Nash and Moser, Bull. Amer. Math. Soc. **7** (1982), 65-222.

[4] H. Hogbe-Nlend, Bornologies and functional analysis, North-Holand Mathematical Studies 62, 1976.

[5] H. Keller, Differential calculus in locally convex spaces. Berlin: Springer-Verlag, 1974.

[6] J. Mawhin and M. Willem, Origin and evolution of the Palais-Smale condition in critical point theory, J. Fixed Point Theory App. **2** (2010) 265-290.

[7] K-H. Neeb, Toward a lie theory of locally convex groups, Jpn. J. Math. **2** (2006), 291-468.

[8] R. S. Palais and S. Smale, A generalized Morse theory, Bull. Amer. Math. Soc. **70** (1964), 165-172.

[9] J. H. Qiu, Ekeland's variational principle in Fréchet spaces and the density of extremal points, Studia Math., **1** (2005), 81-94

[10] T. Wurzacher, Fermionic Second Quantization and the Geometry of the Restricted Grassmannian. Infinite dimensional Kähler manifolds (Oberwolfach, 1995), 287-375, DMV Sem., 31, Birkhäuser, Basel, (2001), 287-375.

Topology dept., Institute of Mathematics of NAS of Ukraine, Tereshchenkivska st. 3, Kyiv, 01601 Ukraine

E-mail address: kaveh@imath.kiev.ua